
QUANTUM MECHANICS NEEDS NO INTERPRETATIONLubomír SKÁLA^{a1,b,*} and Vojtěch KAPSA^{a2}

^a Faculty of Mathematics and Physics, Charles University,
Ke Karlovu 3, 121 16 Prague 2, Czech Republic; e-mail: ¹ skala@karlov.mff.cuni.cz,
² kapsa@karlov.mff.cuni.cz

^b Department of Applied Mathematics, University of Waterloo,
Waterloo, Ontario N2L 3G1, Canada

Received December 21, 2004

Accepted March 29, 2005

Dedicated to Professor Josef Paldus on the occasion of his 70th birthday.

Probabilistic description of results of measurements and its consequences for understanding quantum mechanics are discussed. It is shown that the basic mathematical structure of quantum mechanics like the probability amplitudes, Born rule, probability density current, commutation and uncertainty relations, momentum operator, rules for including scalar and vector potentials and antiparticles can be derived from the definition of the mean values of powers of space coordinates and time. Equations of motion of quantum mechanics, the Klein–Gordon equation, Schrödinger equation and Dirac equation are obtained from the requirement of the relativistic invariance of the theory. The limit case of localized probability densities leads to the Hamilton–Jacobi equation of classical mechanics. Many-particle systems are also discussed.

Keywords: Probability theory; Quantum mechanics; Classical mechanics; Schrödinger equation; Hamilton–Jacobi equation; Probability density current; Equations of motion.

I. INTRODUCTION

Quantum mechanics is one of the most completely tested physical theories (see e.g.^{1–3}). At the same time, the standard approach to introducing quantum mechanics based on the sometimes contrainuitive postulates has rather a mathematical than physical character and exact physical meaning of the postulates and their interpretation is subject of continuing discussion. In this approach, quantum mechanics appears as a field with strange paradoxes and phenomena that are not easy to understand (see e.g.⁴). This is not satisfactory and, in our opinion, it is time (after almost 80 years after formulating the basic principles of quantum mechanics) to replace this approach by a more physical one based on a more direct description of mea-

surements. Only such approach can clarify physical meaning of assumptions made in quantum mechanics. Paraphrasing the title of the recent paper by Fuchs and Peres⁵ we can say: Quantum theory needs no interpretation – it needs derivation from description of measurements. In this sense, our approach can be understood as extension and justification of the standard interpretation of quantum mechanics.

We note that our approach is different from that used usually in physics: To explain experimental results, one introduces some physical quantities and evolution equations for these quantities. Then, consequences of these equations are investigated and compared with experiment. In our approach, we describe results of measurements in a probabilistic way and ask what is the mathematical apparatus that can describe this situation. In this way, the basic mathematical structure of quantum theory except for equations of motion is obtained. Equations of motion are found from the requirement of the relativistic invariance of the probabilistic description.

Probably the best approach is to start with measurement of the space coordinates and time. In this paper, we show that the basic mathematical structure of quantum mechanics like the probability amplitudes, Born rule, probability density current, commutation and uncertainty relations, momentum operator, rules for including scalar and vector potentials and anti-particles can be derived from the definition of the mean values of the space coordinates and time (Sections II–VI). Equations of motion of quantum mechanics, the Klein–Gordon equation, Schrödinger equation and Dirac equation are obtained from the requirement of the relativistic invariance of the theory (Section VII). The limit case of localized probability densities yields the Hamilton–Jacobi equation of classical mechanics (Section VIII). Generalization to many-particle systems is performed in Section IX.

II. BORN RULE

Physical measurements are imperfect and repeated measurements of the same quantity under the same experimental conditions yield different results. The simplest characteristic of such measurements is the mean value of results of repeated measurements and their higher moments. It is the starting point of the following discussion.

Results of repeated measurements of the coordinate x can be characterized by the mean values of x^n

$$\langle x^n \rangle = \int x^n \rho(\mathbf{r}) dV, \quad n = 0, 1, 2, \dots, \quad (1)$$

where integration is carried out over the whole space, $dV = dx dy dz$ and $\rho(\mathbf{r}) \geq 0$ is a normalized probability density

$$\int \rho dV = \langle x^0 \rangle = 1. \quad (2)$$

First, we perform integration by parts with respect to the variable x in Eq. (1) and get

$$\frac{x^{n+1}}{n+1} \rho \Big|_{x=-\infty}^{\infty} - \int \frac{x^{n+1}}{n+1} \frac{\partial \rho}{\partial x} dV = \langle x^n \rangle. \quad (3)$$

Assuming that the first term in this equation equals zero for physically reasonable ρ we get

$$\int x^{n+1} \frac{\partial \rho}{\partial x} dV = -(n+1) \langle x^n \rangle, \quad n = 0, 1, 2, \dots \quad (4)$$

We will show that this simple equation has remarkable consequences.

This equation can be rewritten in form of the inner product

$$(u, v) = -(n+1) \langle x^n \rangle \quad (5)$$

defined in the usual way

$$(u, v) = \int u^*(\mathbf{r}) v(\mathbf{r}) dV. \quad (6)$$

Here, the star denotes the complex conjugate and functions u and v can be taken in general form

$$u = x^{n+1} \psi, \quad (7)$$

$$v = \frac{1}{\psi^*} \frac{\partial \rho}{\partial x}, \quad (8)$$

where $\psi = \psi(\mathbf{r})$ is an arbitrary complex function that will be discussed below. Due to Eq. (5), the Schwarz inequality $(u, u)(v, v) \geq |(u, v)|^2$ can be written in the form

$$(u, u)(v, v) \geq (n+1)^2 \langle x^n \rangle^2, \quad n = 0, 1, 2, \dots, \quad (9)$$

where

$$(u, u) = \int x^{2(n+1)} |\psi|^2 dV, \quad n = 0, 1, 2, \dots \quad (10)$$

and

$$(\nu, \nu) = \int \frac{1}{|\psi|^2} \left(\frac{\partial \rho}{\partial \mathbf{x}} \right)^2 dV. \quad (11)$$

Till now, ψ could be an arbitrary complex function and integrals (u, u) and (ν, ν) had no relation to $\langle x^n \rangle$. Now, we require that inequality (9) does not contain some abstract quantities (u, u) but physically relevant mean values of x^n ($n = 2, 4, 6, \dots$) and suppose

$$\int x^n \rho dV = \int x^n |\psi|^2 dV, \quad n = 2, 4, 6, \dots \quad (12)$$

These conditions do not determine the relation between ρ and $|\psi|^2$ uniquely. For this reason, we assume also that ρ and $|\psi|^2$ have the same norm

$$\int \rho dV = \int |\psi|^2 dV = 1 \quad (13)$$

and the same mean values of odd powers of x

$$\int x^n \rho dV = \int x^n |\psi|^2 dV, \quad n = 1, 3, 5, \dots \quad (14)$$

Then, we obtain relation between the probability density ρ and probability amplitude ψ in form of the Born rule

$$\rho = |\psi|^2 \quad (15)$$

or

$$\psi = \sqrt{\rho} e^{is}, \quad (16)$$

where $s = s(\mathbf{r})$ is an arbitrary real function. Physical meaning of the phase s will be discussed later.

We note that for $\rho = |\psi|^2$ the integral (ν, ν) is the so-called Fisher information⁶⁻¹⁰.

The Born rule $\rho = |\psi|^2$ is the only relation between ρ and ψ for which $\int x^n \rho dV = \int x^n |\psi|^2 dV$, $n = 0, 1, 2, \dots$. Significance of the Born rule follows from the fact that inequalities (9) contain for $\rho = |\psi|^2$ physically relevant quantities $\int x^n \rho dV$, $n = 2, 4, 6, \dots$. For other relations between ρ and ψ , this physically important property is in general lost.

III. PROBABILITY DENSITY CURRENT

We note that to describe physical systems, we have to specify not only the space-time probability distribution ρ , but also its evolution in space-time. This information can be encoded into the complex part of the probability amplitude ψ . Here, we can proceed similarly as in continuum mechanics, where not only the density ρ but also the density current

$$\mathbf{j}_k = \rho \mathbf{v}_k, \quad k = 1, 2, 3 \quad (17)$$

related to components of the velocity \mathbf{v}_k is introduced. Writing the "velocity" in the form

$$\mathbf{v}_k = \frac{\partial s}{\partial \mathbf{x}^k}, \quad (18)$$

where $s = s(\mathbf{r}, t)$ is a real function we get

$$\mathbf{j}_k = \rho \frac{\partial s}{\partial \mathbf{x}^k} = \sqrt{\rho} e^{-is} \sqrt{\rho} e^{is} \frac{\partial s}{\partial \mathbf{x}^k} = \sqrt{\rho} e^{-is} (-i) \frac{\partial(\sqrt{\rho} e^{is})}{\partial \mathbf{x}^k} + \frac{i}{2} \frac{\partial \rho}{\partial \mathbf{x}^k}. \quad (19)$$

Using the complex probability amplitude (16) we have

$$\mathbf{j}_k = \psi^* \left(-i \frac{\partial \psi}{\partial \mathbf{x}^k} \right) + \frac{i}{2} \frac{\partial \rho}{\partial \mathbf{x}^k}. \quad (20)$$

However, the probability density current has to be real. Calculating the real part of \mathbf{j}_k we obtain the final expression

$$\mathbf{j}_k = \frac{1}{2} \left[\psi^* \left(-i \frac{\partial \psi}{\partial \mathbf{x}^k} \right) + c. c. \right] = \frac{1}{2i} \left(\psi^* \frac{\partial \psi}{\partial \mathbf{x}^k} - \psi \frac{\partial \psi^*}{\partial \mathbf{x}^k} \right). \quad (21)$$

Except for a multiplicative factor, this formula agrees with the expression for the probability density current known from quantum mechanics. Complex probability amplitudes ψ are necessary to obtain nonzero \mathbf{j}_k . The probability density current depends on the operator $-i(\partial/\partial \mathbf{x}^k)$. Except for the factor \hbar , this operator agrees with the momentum operator $\hat{p}_k = -i\hbar(\partial/\partial \mathbf{x})$ known from quantum mechanics. In agreement with rules of quantum mechanics, the probability amplitudes ψ and $\psi \exp(i\alpha)$, where α is a real constant, yield the same probability density ρ and probability density current \mathbf{j}_k .

IV. COMMUTATION RELATION

Now we return back to Eq. (4) for $n = 0$. Using the Born rule (15) and Eq. (2) we rewrite Eq. (4) in form containing the probability amplitude ψ

$$\int x \left(\frac{\partial \psi^*}{\partial x} \psi + \psi^* \frac{\partial \psi}{\partial x} \right) dV = -1. \quad (22)$$

Multiplying this equation by $-i$ we obtain the equation

$$\int \left[(x\psi)^* \left(-i \frac{\partial \psi}{\partial x} \right) - \left(-i \frac{\partial \psi}{\partial x} \right)^* x\psi \right] dV = i \quad (23)$$

or in the operator form

$$[x, -i(\partial/\partial x)] = i. \quad (24)$$

This commutation relation is straightforward consequence of Eq. (4) for $n = 0$ and the Born rule (15) and need not be postulated. Except for the factor \hbar determining the choice of units, this commutation relation agrees with the commutation relation between the coordinate and momentum operator known from quantum mechanics.

V. UNCERTAINTY RELATIONS

The uncertainty relation for the coordinate x and the operator $-i(\partial/\partial x)$ can be derived in a standard way from the commutation relation (24) or, equivalently, by means of a simple calculation of (v, v)

$$(v, v) = \int \frac{1}{|\psi|^2} \left(\frac{\partial \rho}{\partial x} \right)^2 dV = 4 \int \frac{1}{|\psi|^2} \left[\operatorname{Re} \left(\psi^* \frac{\partial \psi}{\partial x} \right) \right]^2 dV \leq 4 \int \left| -i \frac{\partial \psi}{\partial x} \right|^2 dV. \quad (25)$$

Substituting expressions (10) and (25) into inequality (9) for $n = 0$ we get, except for \hbar^2 , the uncertainty relation in the form known from quantum mechanics

$$\int x^2 |\psi|^2 dV \int \left| -i \frac{\partial \psi}{\partial x} \right|^2 dV \geq \frac{1}{4}. \quad (26)$$

This result can be further generalized. Using integration by parts and the condition $\rho \rightarrow 0$ for $x \rightarrow \pm\infty$, Eq. (22) can be generalized as

$$\int [(x-a)\psi]^* \left[\frac{\partial \psi}{\partial x} - i b \psi \right] dV + \int \left[\frac{\partial \psi}{\partial x} - i b \psi \right]^* [(x-a)\psi] dV = -1, \quad (27)$$

where a and b are real constants. From this equation, a more general form of the uncertainty relation can be obtained

$$\int (x-a)^2 |\psi|^2 dV \int |-i\partial\psi/\partial x - b\psi|^2 dV \geq \frac{1}{4}. \quad (28)$$

Minimum of the left-hand side is obtained for

$$a = \int \psi^* x \psi dV = \langle x \rangle \quad (29)$$

and

$$b = \int \psi^* (-i\partial\psi / \partial x) dV = \langle -i\partial/\partial x \rangle. \quad (30)$$

Except for the factor \hbar , the resulting uncertainty relation with a and b given by the last two equations agrees with the well-known Heisenberg uncertainty relation. Again, it can be obtained from Eq. (4) for $n = 0$ and the Born rule (15). Another general discussion of uncertainty relations can be found in¹¹.

Uncertainty relations are a general consequence of Eq. (1) and must appear in any probabilistic theory of this kind, including quantum mechanics. There are two important quantities appearing in the uncertainty relations: the coordinate x and the operator $-i(\partial/\partial x)$. Similar quantities, namely the coordinate x and the momentum operator $\hat{p}_k = -i\hbar(\partial/\partial x)$ appear also in quantum mechanics.

It is worth noting that Eq. (27) remains valid also in the case if b is a real function $b = f_x(\mathbf{r}, t)$. This means that the operator $-i(\partial\psi/\partial x)$ can be replaced by the operator $-i(\partial\psi/\partial x) - f_x$ and the commutation relation (24) and the uncertainty relation (28) can be further generalized. Therefore, general structure of the probability theory remains preserved for any real function f_x . In physics, different functions f_x , f_y and f_z correspond to different components of the electromagnetic vector potential $\mathbf{A} = (A_x, A_y, A_z)$ multiplied by the charge q of the particle. Except for \hbar and q , it agrees with the rule $-i\hbar\nabla \rightarrow -i\hbar\nabla - q\mathbf{A}$ for including the vector potential \mathbf{A} into quantum theory (for charge, see the end of Section VI). We note also that the kinetic energy in quantum mechanics $T = (\hbar/2m) \int |\nabla\psi|^2 dV$ equals the space Fisher information $\int |\nabla\rho|^2 / \rho dV$ multiplied by $\hbar^2 / (8m)$.

VI. TIME

Time can be discussed analogously as the space coordinates, however, there are some important differences that has to be taken into consideration.

Assuming that there are given initial conditions for $\psi(\mathbf{r}, t = 0)$ the probability amplitude $\psi(\mathbf{r}, t)$, $t > 0$ gives the probability description of measurements at later times. Therefore, time evolution has unidirectional character from given initial conditions to the relative probability of results of (yet unperformed) measurements at later times. If this measurement is actually performed, the probabilistic description must be replaced by a specific result obtained from the performed measurement. This is the basis of two different evolution schemes in quantum mechanics: time evolution described by the evolution equation like the Schrödinger equation and reduction or collapse of the wave function. In this paper, we are interested in the former case. Detailed description of the reduction of the probability amplitude is not needed in our approach.

In standard quantum mechanics, the probability amplitudes obey the normalization condition $\int |\nabla\psi|^2 dV = 1$ valid at all times and the integral over time $\int_0^\infty \int |\nabla\psi|^2 dV dt$ goes to infinity. This situation can be compared with that for a free particle. For a free particle, the integral $\int |\psi|^2 dV$ goes to infinity and ψ is usually normalized by means of the Dirac δ -function. For time, similar approach cannot be used for two reasons. First, we do not perform here integration over all times, but from the initial condition at $t = 0$ to infinity. Second, if the integral $\int_0^\infty \int |\nabla\psi|^2 dV dt$ goes to infinity, we cannot define the mean time by analogy with Eq. (1) and proceed similarly as in the preceding Sections. For these reasons, we assume that not only the integral $\int |\psi|^2 dV$ but also the integral

$$\int_0^\infty \int |\psi|^2 dV dt = 1 \quad (31)$$

equals one and proceed by analogy with the space coordinates. In this way, we get the operator $-i(\partial/\partial t)$, obtain the corresponding commutation and uncertainty relations and introduce the scalar potential. At the end of our discussion, we will assume that $\int |\psi|^2 dV$ changes very slowly in time, normalize the probability amplitude by means of the usual condition $\int |\psi|^2 dV = 1$ and perform transition to standard quantum mechanics.

First, we define the time component of the probability density current by the equation analogous to Eqs. (17) and (18)

$$j_t = -\rho \frac{\partial s}{\partial t} \quad (32)$$

and obtain expression similar to Eq. (21)

$$j_t = \frac{1}{2} \left[\psi^* \left(i \frac{\partial \psi}{\partial t} \right) + c. c. \right]. \quad (33)$$

Except for a factor, this quantity equals the time component of the probability density current $j_0 = \text{Re}[\psi^* i \hbar (\partial \psi / \partial x^0)] / m_0$ known from relativistic quantum mechanics, where $x^0 = ct$ and m_0 is the rest mass. Then, by analogy with Eq. (23) we derive the equation

$$\int_{t=0}^{\infty} \int \left[\left(i \frac{\partial \psi}{\partial t} \right)^* t \psi - (t \psi)^* \left(i \frac{\partial \psi}{\partial t} \right) \right] dV dt = i. \quad (34)$$

One can introduce also a real constant d into this equation

$$\int_{t=0}^{\infty} \int \left[\left(i \frac{\partial \psi}{\partial t} - d \psi \right)^* t \psi - (t \psi)^* \left(i \frac{\partial \psi}{\partial t} - d \psi \right) \right] dV dt = i. \quad (35)$$

The uncertainty relation for time can be written in form analogous to Eq. (28)

$$\int_0^{\infty} \int t^2 |\psi|^2 dV dt \int_0^{\infty} \int |i \partial \psi / \partial t - d \psi|^2 dV dt \geq \frac{1}{4}. \quad (36)$$

Minimum of the left-hand side is obtained for

$$d = \frac{1}{2} \left[\int_0^{\infty} \int \psi^* i (\partial y / \partial t) dV dt + c. c. \right]. \quad (37)$$

Equation (36) is valid also if d is replaced by a real function $f_0(\mathbf{r}, t)$.

To illustrate meaning of Eq. (36) we assume decaying probability amplitude with the life time $\tau > 0$

$$\psi(\mathbf{r}, t) = \frac{1}{\sqrt{\tau}} e^{-i\omega t - t/(2\tau)} \psi(\mathbf{r}), \quad (38)$$

where the space part of the probability amplitude is normalized by the usual condition $\int |\psi(\mathbf{r})|^2 dV = 1$. In this case, we get from Eqs. (36) and (37)

$$\int_0^{\infty} \int t^2 |\psi|^2 dV dt = 2\tau^2, \quad (39)$$

$d = \omega$ and

$$\int_0^\infty \int |\mathrm{i}\partial\psi/\partial t - d\psi|^2 \mathrm{d}V \mathrm{d}t = \frac{1}{4\tau^2}. \quad (40)$$

Therefore, uncertainty relation (36) gives the relation between the mean square time $\langle t^2 \rangle = 2\tau^2$ and the square of the imaginary part of the complex frequency $\omega - \mathrm{i}/(2\tau)$ and has the meaning of the time-energy uncertainty relation.

In agreement with our understanding of direction of time, we assume that direct physical meaning have only the probability amplitudes corresponding to the non-negative values of the time component of the probability density current integrated over the whole space

$$\int j_t \mathrm{d}V = -\int \rho \frac{\partial s}{\partial t} \mathrm{d}V \geq 0. \quad (41)$$

If this quantity is negative, its sign can be reversed by the transformation $\psi \rightarrow \psi^*$ changing the sign of the phase s and the probability density currents j_k and j_t . Performing this transformation we get from Eq. (35) for $d = f_0$

$$\int_{t=0}^\infty \int \left[\left(\mathrm{i} \frac{\partial \psi}{\partial t} + f_0 \psi \right)^* t \psi - (t \psi)^* \left(\mathrm{i} \frac{\partial \psi}{\partial t} + f_0 \psi \right) \right] \mathrm{d}V \mathrm{d}t = \mathrm{i} \quad (42)$$

and see that this transformation changes the sign of f_0 .

Similar discussion can be done also for the space coordinates. As a result, the transformation $\psi \rightarrow \psi^*$ leads to change of sign of the functions f_0 and f_k , $k = 1, 2, 3$ that can for example be respected by putting $f_0 = qU$ and $f_k = qA_k$, where U and A_k are the scalar and vector electromagnetic potentials. Therefore, the probability amplitudes ψ and ψ^* describe particles that differ by the sign of their charge and general structure of the probability theory and unidirectional character of time lead to the existence of particles and antiparticles.

Except for \hbar , our results agree with the rules $\mathrm{i}\hbar(\partial/\partial t) \rightarrow \mathrm{i}\hbar(\partial/\partial t) - qU$ and $-\mathrm{i}\hbar\nabla \rightarrow -\mathrm{i}\hbar\nabla - q\mathbf{A}$ for including the electromagnetic potentials into quantum theory. These potentials representing different physical scenarios do not appear among the variables of the probability amplitude and describe non-quantized classical fields.

Now, we perform transition to standard quantum mechanics. In this limit case the integration over time need not be performed and the proba-

bility density can be normalized over the space only $\int |\psi|^2 dV = 1$. At the same time, the uncertainty relation (36) loses its original meaning and time becomes a parameter rather than a dynamical variable. It is the first reason for a different role of time and space coordinates in quantum mechanics. The second reason is that the operator $i(\partial/\partial t)$ appears in equations of motion like the Schrödinger equation and does not represent an independent physical quantity.

It is worth noting that to obtain results of Sections II–VI no evolution equation has been needed. Therefore, this part of the mathematical formalism of quantum mechanics follows directly from the probabilistic description of results of measurements. It is also interesting that the Planck constant \hbar does not appear in our discussion and can be included by multiplying Eqs. (23) and (34) by \hbar . Therefore, the Planck constant determines the units used in measurements and scales at which the probabilistic character of measurements is important.

VII. EQUATIONS OF MOTION

To find equations of motion we require relativistic invariance of the theory. In this respect, our approach is different from that based on the principle of extreme physical information or minimum of the Fisher information^{8,12}.

First we note that all quantities discussed above depend on ψ or its first derivatives with respect to time and space coordinates. Returning to our scheme used in Section VI we can create real relativistic invariant from the first derivatives of ψ appearing in the uncertainty relations (28) and (36) for $a = b = d = 0$

$$\int_0^\infty \int \left(\frac{1}{c^2} \left| \frac{\partial \psi}{\partial t} \right|^2 - \sum_{k=1}^3 \left| \frac{\partial \psi}{\partial x^k} \right|^2 \right) dV dt = \text{const}, \quad (43)$$

where c is the speed of light.

Integral $\int_0^\infty \int |\partial\psi/\partial t|^2 dV dt$ has meaning of the time Fisher information and is non-negative. Similar conclusion applies also to $\int_0^\infty \int |\partial\psi/\partial x^k|^2 dV dt$, $k = 1, 2, 3$. However, since Eq. (43) must be valid in all cases including the case $\partial\psi/\partial x^k = 0$ (in the language of quantum mechanics, it corresponds to zero momentum and zero kinetic energy) we can conclude that $\text{const} \geq 0$.

In this equation, we can perform integration by parts with respect to all variables. For example, we get for time

$$\int_0^\infty \int \frac{\partial \psi^*}{\partial t} \frac{\partial \psi}{\partial t} dV dt = \frac{1}{2} \left[\int \left(\psi^* \frac{\partial \psi}{\partial t} + c.c. \right) dV \right]_0^\infty - \frac{1}{2} \int_0^\infty \int \left(\psi^* \frac{\partial^2 \psi}{\partial t^2} + c.c. \right) dV dt. \quad (44)$$

However, the first integral on the right-hand side can be expressed as $\partial(\int |\psi|^2 dV)/\partial t$ and disappears in the limit of standard quantum mechanics when $\int |\psi|^2 dV = 1$. An analogous result can be obtained also for the variables x^k assuming that $\partial|\psi|^2/\partial x^k$ for $x^k \rightarrow -\infty$ and $x^k \rightarrow \infty$ equal. In standard quantum mechanics, this condition is obeyed for a free particle as well as for the bound states.

Now, we perform transition to standard quantum mechanics with the wave function normalized in the usual way and get equation that has to be valid for all ψ

$$\frac{1}{2} \int \left[\psi^* \left(\Delta - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \text{const} \right) \psi + c. c. \right] dV = 0. \quad (45)$$

This condition is obeyed if the probability amplitudes fulfill the equation

$$\left(\Delta - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \text{const} \right) \psi = 0. \quad (46)$$

Since $\text{const} \geq 0$, we can put $\text{const} = m_0^2 c^2 / \hbar^2$, where m_0 is another constant, known as the rest mass of the particle. Therefore, requirement of the relativistic invariance applied to quantities appearing in the probabilistic formulation leads to the Klein-Gordon equation for a free particle.

The non-relativistic time Schrödinger equation can be obtained from the Klein-Gordon equation by using the transformation

$$\psi = e^{m_0 c^2 t / (\hbar)} \varphi, \quad (47)$$

where φ is the probability amplitude appearing in the Schrödinger equation. This transition is well-known and will not be discussed here (see e.g.¹³).

The Dirac equation can be derived by replacing the probability amplitude ψ in Eq. (43) by a column vector with four components

$$\int_0^\infty \int \frac{1}{c^2} \left(\frac{\partial \psi^+}{\partial t} \frac{\partial \psi}{\partial t} - \sum_{k=1}^3 \frac{\partial \psi^+}{\partial x^k} \frac{\partial \psi}{\partial x^k} \right) dV dt = \text{const}, \quad (48)$$

where the cross denotes the hermitian conjugate. Inserting the γ^μ matrices¹³ into this equation, putting $\text{const} = m_0^2 c^2 / \hbar^2$ and using Eq. (31) we get

$$\int_0^\infty \int \left[\frac{1}{c^2} \left(\gamma^0 \frac{\partial \psi}{\partial t} \right)^+ \left(\gamma^0 \frac{\partial \psi}{\partial t} \right) - \sum_{k=1}^3 \left(\gamma^k \frac{\partial \psi}{\partial x^k} \right)^+ \left(\gamma^k \frac{\partial \psi}{\partial x^k} \right) - \frac{m_0^2 c^2}{\hbar^2} \psi^+ \psi \right] dV dt = 0. \quad (49)$$

Then, using properties of the γ^μ matrices and assuming that the integration by parts can be used analogously as in case of Eq. (44) the last equation leads in the limit of standard quantum mechanics to (see also^{7,8})

$$\int \left(\frac{\gamma^0}{c} \frac{\partial \Psi}{\partial t} - \sum_{k=1}^3 \gamma^k \frac{\partial \Psi}{\partial x^k} - \frac{i m_0 c}{\hbar} \Psi \right)^+ \left(\frac{\gamma^0}{c} \frac{\partial \Psi}{\partial t} + \sum_{k=1}^3 \gamma^k \frac{\partial \Psi}{\partial x^k} + \frac{i m_0 c}{\hbar} \Psi \right) dV = 0. \quad (50)$$

The operator in the first parentheses is the hermitian conjugate of that in the second ones. Assuming that the expression in the second parentheses equals zero, we obtain the Dirac equation for a free particle

$$\frac{\gamma^0}{c} \frac{\partial \Psi}{\partial t} + \sum_{k=1}^3 \gamma^k \frac{\partial \Psi}{\partial x^k} + \frac{i m_0 c}{\hbar} \Psi = 0. \quad (51)$$

We can see that requirement of the relativistic invariance of the probabilistic description yields all the basic equations of motion of quantum mechanics. The scalar and vector potentials can be included by means of the rules $i\hbar(\partial/\partial t) \rightarrow i\hbar(\partial/\partial t) - qU$ and $-i\hbar\nabla \rightarrow -i\hbar\nabla - q\mathbf{A}$ discussed above.

VIII. CLASSICAL MECHANICS

To derive the Hamilton–Jacobi equation for a free particle we proceed as follows. The probability amplitude is assumed in the form

$$\Psi = e^{is/\hbar} = e^{is_1/\hbar} e^{-s_2/\hbar}, \quad (52)$$

where s_1 and s_2 are the real and imaginary parts of s , respectively. In the limit of standard quantum mechanics mentioned above, Eq. (43) with $\text{const} = m_0^2 c^2 / \hbar^2$ can be replaced by the equation

$$\frac{1}{c^2} \int \left| \frac{\partial s}{\partial t} \right|^2 |\Psi|^2 dV = \int |\nabla s|^2 |\Psi|^2 dV + m_0^2 c^2. \quad (53)$$

Now we assume that the probability density

$$\rho = |\Psi|^2 = e^{-2s_2/\hbar} \quad (54)$$

has very small values everywhere except for the vicinity of the point $\langle \mathbf{r} \rangle$, where it achieves its maximum and the first derivatives of s_2 at this point equal zero

$$\left. \frac{\partial s_2}{\partial x^k} \right|_{\mathbf{r}=\langle \mathbf{r} \rangle} = 0, \quad k = 1, 2, 3. \quad (55)$$

In such a case, the probability density can be replaced by the δ -function

$$|\psi|^2 = \delta(\mathbf{r} - \langle \mathbf{r} \rangle) \quad (56)$$

and probabilistic character of the theory disappears. Equations (53)–(56) then lead to the relativistic equation

$$\frac{1}{c^2} \left(\frac{\partial s_1(\langle \mathbf{r} \rangle, \langle t \rangle)}{\partial t} \right)^2 = [\nabla s_1(\langle \mathbf{r} \rangle, \langle t \rangle)]^2 + m_0^2 c^2. \quad (57)$$

We note that Eq. (56) corresponds to the limit $\hbar \rightarrow 0$ in Eq. (54). Therefore, s_1 in Eq. (57) is in fact the first term of the expansion of s_1 into the power series in \hbar

$$s_1 = s_1|_{\hbar=0} + \dots \quad (58)$$

Further, we replace the mean coordinates $\langle \mathbf{r} \rangle$ by \mathbf{r} as it is usual in classical mechanics and introduce the classical non-relativistic action $S(\mathbf{r}, t)$

$$s_1 = S - m_0 c^2 t. \quad (59)$$

Equation (57) then leads to

$$\frac{1}{c^2} \left(\frac{\partial S}{\partial t} - m_0 c^2 \right)^2 = (\nabla S)^2 + m_0^2 c^2. \quad (60)$$

In the non-relativistic limit $|\partial S/\partial t| \ll m_0 c^2$, the last equation yields the Hamilton–Jacobi equation for a free particle

$$\frac{\partial S}{\partial t} + \frac{(\nabla S)^2}{2m_0} = 0. \quad (61)$$

Thus, the Hamilton–Jacobi equation can be obtained from the probabilistic description of results of measurements in the limit of δ -like probability densities and non-relativistic approximation. The scalar and vector potentials U and \mathbf{A} can be included by means of the rules $\partial S/\partial t \rightarrow \partial S/\partial t + qU$ and

$\nabla S \rightarrow \nabla S - q\mathbf{A}$ following from the rules $i\hbar(\partial/\partial t) \rightarrow i\hbar(\partial/\partial t) - qU$ and $-i\hbar\nabla \rightarrow -i\hbar\nabla - q\mathbf{A}$ discussed above.

IX. MANY-PARTICLE SYSTEMS

In general, many-particle systems have to be described by quantum field theory. However, if we limit ourselves to quantum mechanics, we can proceed as follows.

The starting point of discussion of the N -particle system is the definition analogous to Eq. (1)

$$\langle \mathbf{r}_j \rangle = \int \mathbf{r}_j \rho(\mathbf{r}_1, \dots, \mathbf{r}_N, t) dV_1 \dots dV_N, \quad j = 1, \dots, N, \quad (62)$$

where ρ is the many particle probability density and \mathbf{r}_j are the coordinates of the j -th particle. Then, discussion can be performed analogously to that given above and the probability amplitude, uncertainty and commutation relations, momentum operators and density currents for all particles can be introduced. The scalar and vector potentials $U(\mathbf{r}_1, \dots, \mathbf{r}_N, t)$ and $\mathbf{A}(\mathbf{r}_1, \dots, \mathbf{r}_N, t)$ and antiparticles can be also discussed.

Equations of motion for N free particles can be found from generalization of the relativistic invariant (43)

$$\int_0^\infty \int \left(\frac{1}{c^2} \left| \frac{\partial \psi}{\partial t} \right|^2 - \sum_{j=1}^N |\nabla_j \psi|^2 \right) dV_1 \dots dV_N dt = \sum_{j=1}^N \frac{m_j^2 c^2}{\hbar^2}, \quad (63)$$

where $\psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t)$ is the N -particle probability amplitude and m_j denotes the rest mass of the particle.

Using similar approach as above, we can then obtain the N -particle Schrödinger equation

$$-\sum_{j=1}^N \frac{\hbar^2}{2m_j} \Delta_j \psi = i\hbar \frac{\partial \psi}{\partial t} \quad (64)$$

and the Hamilton-Jacobi equation

$$\frac{\partial S}{\partial t} + \sum_{j=1}^N \frac{(\nabla_j S)^2}{2m_j} = 0. \quad (65)$$

For a system of identical particles, the probability density ρ must be symmetric with respect to the exchange of any two particles i and j . Hence, the

probability amplitude ψ must be symmetric or antisymmetric with respect to such exchanges.

Non-locality of quantum mechanics is related to the many-particle character of the probability density ρ and the corresponding probability amplitude ψ .

It can be seen that probabilistic description of measurements and its relativistic invariance yields also the basic mathematical structure of the many-particle quantum mechanics.

X. CONCLUSIONS

In this paper, we have shown that the basic mathematical structure of quantum mechanics can be derived from the probabilistic description of the results of measurement of the space coordinates and time. Equations of motion of quantum mechanics have been obtained from the requirement of the relativistic invariance of the theory. As the limit case, this approach yields also the Hamilton–Jacobi equation of classical mechanics.

Unperformed experiments have no results. Therefore, it follows from our approach that quantum mechanics does not speak of events in the measured system, but only of results of measurements, implying the existence of external measuring apparatus.

Since our approach makes it possible to obtain the most significant parts of the mathematical formalism of quantum mechanics from the probabilistic description of results of measurements, we believe that it is a natural and physically satisfactory starting point to understanding this field. It contributes also to understanding quantum theory as correctly formulated probabilistic description of measurements that can describe physical phenomena at different levels of accuracy from the simplest models to very complex ones.

This work was supported by the Grant Agency of the Czech Republic (grant No. 202/03/0799) and the Ministry of Education, Youth and Sports of the Czech Republic (grant No. 0021620835).

REFERENCES

1. Zeilinger A.: *Rev. Mod. Phys.* **1999**, *71*, S288.
2. Arndt M., Nairz O., Zeilinger A. in: *Quantum [Un]speakables. From Bell to Quantum Information* (R. A. Bertlmann and A. Zeilinger, Eds). Springer, Berlin 2002.
3. Scully M. O., Englert B. G., Walther H.: *Nature* **1991**, *351*, 111.
4. Laloë F.: *Am. J. Phys.* **2001**, *69*, 655.
5. Fuchs Ch. A., Peres A.: *Physics Today* **2000**, 70.

6. Fisher R. A.: *Proc. Cambr. Phil. Soc.* **1925**, 22, 700.
7. Roy Frieden B., Soffer B. H.: *Phys. Rev. E: Stat. Phys., Plasmas, Fluids, Relat. Interdiscip. Top.* **1995**, 52, 2274.
8. Roy Frieden B.: *Physics from Fisher Information*. Cambridge University Press, Cambridge 1998.
9. Cover T., Thomas J.: *Elements of Information Theory*. Wiley, New York 1991.
10. Hall M. J. W.: *Phys. Rev. A: At., Mol., Opt. Phys.* **2000**, 62, 012107.
11. Roy Frieden B.: *Phys. Lett. A* **1992**, 169, 123.
12. Hall M. J. W.: *Phys. Rev. A: At., Mol., Opt. Phys.* **2001**, 64, 052103.
13. Reginatto M.: *Phys. Rev. A: At., Mol., Opt. Phys.* **1998**, 58, 1775.
14. Davydov A. S.: *Quantum Mechanics*. Pergamon Press, New York 1976.