

# Large-order behavior of the convergent perturbation theory for anharmonic oscillators

L. Skála,<sup>1,2</sup> J. Čížek,<sup>2,1</sup> E. J. Weniger,<sup>3,1,2</sup> and J. Zamastil<sup>1</sup>

<sup>1</sup>Faculty of Mathematics and Physics, Charles University, Ke Karlovu 3, 12116 Prague 2, Czech Republic

<sup>2</sup>Department of Applied Mathematics, University of Waterloo, Waterloo, Ontario, Canada N2L 3G1

<sup>3</sup>Institut für Physikalische und Theoretische Chemie, Universität Regensburg, D-93040 Regensburg, Germany

(Received 28 January 1998; revised manuscript received 6 July 1998)

Using the large-order formula for the coefficients of the divergent weak-coupling series for the energy of the anharmonic oscillators, we derive a simple analytic large-order formula for the coefficients of the convergent renormalized strong-coupling series. This formula is valid for all the states of the anharmonic oscillators defined by the Hamiltonians  $H = p^2 + x^2 + \beta x^{2m}$  with  $m \geq 2$ . A further generalization of this formula is also proposed. Numerical tests of the formula are performed for the quartic, sextic, octic, and decadic oscillator with the help of asymptotic analysis. Further it is shown that the renormalized strong-coupling perturbation expansion converges for all the states of these oscillators and for all physically relevant  $\beta \in [0, \infty)$ .

[S1050-2947(99)04901-X]

PACS number(s): 03.65.-w, 02.30.Lt, 02.70.-c

## I. INTRODUCTION

We investigate the Schrödinger equation  $H\psi = E(\beta)\psi$  for the anharmonic oscillators, where

$$H = p^2 + x^2 + \beta x^{2m}, \quad \beta \geq 0, \quad m \geq 2. \quad (1)$$

As is well known,  $E(\beta)$  can be expressed as a *weak-coupling* perturbation series in powers of  $\beta$ ,

$$E(\beta) = \sum_{n=0}^{\infty} b_n \beta^n, \quad (2)$$

which diverges for every  $\beta > 0$  [1–6]. The large-order behavior of the coefficients  $b_n$  follows from [3]

$$b_n = (-1)^{n+1} \frac{(m-1)2^K}{\pi^{3/2} K! 2^{2n-1}} \Gamma(n(m-1) + K + 1/2) a^{n(m-1) + K + 1/2}, \quad K \geq 0, \quad (3)$$

$$a = \frac{\Gamma(2m/(m-1))}{\Gamma^2(m/(m-1))}, \quad (4)$$

where  $K$  is the index of the excitation.

With the help of the scaling transformation  $x \rightarrow \beta^{-1/[2(m+1)]}x$ ,  $H$  can be expressed as [2]

$$H = \beta^{1/(m+1)} [p^2 + \beta^{-2/(m+1)} x^2 + x^{2m}]. \quad (5)$$

Consequently,  $E(\beta)$  also possesses the *strong-coupling* expansion

$$E(\beta) = \beta^{1/(m+1)} \sum_{n=0}^{\infty} K_n \beta^{-2n/(m+1)}, \quad (6)$$

which converges if  $\beta$  is sufficiently large [2].

Alternative perturbative approaches based upon renormalization (Wick ordering [7] or scaling [7–11]) have considerable conceptual and technical advantages. In the quartic case,

Wick ordering and scaling are closely related and they differ by a numerical factor in the effective coupling constant. In this paper we do scaling according to  $x \rightarrow \sqrt{\tau}x$ , where  $\tau$  and  $\beta$  are related by the  $m$ -dependent equation

$$\beta = (1 - \tau^2) / (B_m \tau^{m+1}), \quad (7)$$

with  $B_m = m(2m-1)!!/2^{m-1}$  [9]. This transformation maps the physically relevant *unbounded* interval  $\beta \in [0, \infty)$  onto the *bounded* interval  $\tau \in [1, 0)$ . With the help of Eq. (7), the Hamiltonian (1) can be expressed in terms of a renormalized Hamiltonian  $H_R$  [9,10],

$$H = H_R / \tau, \quad (8)$$

$$H_R(\tau) = p^2 + x^2 + (1 - \tau^2)(x^{2m}/B_m - x^2) \quad (9)$$

$$= p^2 + x^{2m}/B_m + \tau^2(x^2 - x^{2m}/B_m). \quad (10)$$

The *renormalized energy*

$$E_R(\tau) = \tau E(\beta) \quad (11)$$

can either be expressed as a divergent *renormalized weak-coupling* expansion in  $1 - \tau^2$  [10],

$$E_R(\tau) = \sum_{n=0}^{\infty} c_n (1 - \tau^2)^n, \quad (12)$$

or as a *renormalized strong-coupling* expansion in  $\tau^2$  [11],

$$E_R(\tau) = \sum_{n=0}^{\infty} \Gamma_n \tau^{2n}. \quad (13)$$

The weak-coupling expansion (12) diverges almost as strongly as the weak-coupling expansion (2) [7,10]. However, the strong-coupling expansion (13) has some very useful properties [11,12].

For the ground and first excited states of the quartic anharmonic oscillator, we computed numerically 200 coeffi-

coefficients  $\Gamma_n$  with high accuracy [12]. From these data, we obtained the following large-order behavior:

$$\Gamma_n = A^{(K)} (2n)^{(K-1)/2} e^{-2\sqrt{2}n} \left( 1 + \sum_{\nu=1}^{\infty} \frac{a_{\nu}^{(K)}}{(2n)^{\nu/2}} \right). \quad (14)$$

The leading coefficient

$$A^{(K)} = -\frac{12^K}{K!} \frac{4\sqrt{6}}{\pi e^2} \quad (15)$$

was determined from the summation rules for  $\Gamma_n$ . In addition, a few coefficients  $a_{\nu}^{(K)}$  were also determined analytically. The leading term of the large-order formula (14) shows that the renormalized strong-coupling expansion (13) converges for all  $\tau \in [0,1)$  [12].

Our results for the quartic anharmonic oscillator and further numerical results for the sextic and octic anharmonic oscillators [11] indicate that the renormalized strong-coupling expansion (13) actually converges for arbitrary  $m \geq 2, K \geq 0$ , and  $\tau \in [0,1)$ .

The main purpose of this paper is to investigate *analytically* the large-order behavior of the renormalized strong-coupling coefficients  $\Gamma_n$  of general anharmonic oscillators with Hamiltonian (1). We show in Sec. II that their large-order behavior is described by a simple analytic formula which is a generalization of the leading term in Eq. (14) to arbitrary  $m \geq 2$ . In contrast to [12], where Eq. (14) was conjectured from numerical analysis, we use here an analytic approach. We propose also further generalization of the series (14). In Sec. III we compare the large-order formula with the actual values of the  $\Gamma_n$  coefficients and test the validity of the summation rule for  $\Gamma_n$ . These numerical results and the large-order formula for  $\Gamma_n$  show that the renormalized strong-coupling expansion (13) converges for all  $\tau \in [0,1)$ ,  $m \geq 2$ , and  $K \geq 0$ .

Finally, let us mention that our final goal is the resummation of renormalized series both for the case of oscillators ( $\tau \in [0,1)$ ) and double wells [ $\tau \in (-\infty, 0)$ ]. Consideration of the large-order behavior of these coefficients is a step in this program.

## II. LARGE-ORDER FORMULA

For the determination of the large-order behavior of  $\Gamma_n$ , we start from Eq. (3) for the coefficients  $b_n$ . For large  $n$ , it follows from [7,10] that  $c_n = b_n / [e^3 (B_m)^n]$  for  $m=2$ , and  $c_n = b_n / (B_m)^n$  for  $m \geq 3$ . This leads to the large-order behavior valid for  $m \geq 3$ ,

$$c_n = (-1)^{n+1} C \Gamma((m-1)n + K + 1/2) / D^n, \quad (16)$$

$$C = \frac{(m-1)}{\pi^{3/2}} \frac{2^{K+1}}{K!} a^{K+1/2}, \quad (17)$$

$$D = 4B_m a^{1-m}. \quad (18)$$

For  $m=2$ , the constant  $C$  in Eq. (17) has to be divided by  $e^3$ .

For the derivation of the large-order formula for  $\Gamma_n$  we use Eq. (16). First we substitute Eq. (16) into the remainder of the series (12),

$$\begin{aligned} \Delta E_R(\kappa) &= E_R(\kappa) - \sum_{\nu=0}^{n-1} c_{\nu} \kappa^{\nu} \\ &= C \sum_{\nu=n}^{\infty} (-1)^{\nu+1} \left( \frac{\kappa}{D} \right)^{\nu} \Gamma((m-1)\nu + K + 1/2), \end{aligned} \quad (19)$$

where  $\kappa = 1 - \tau^2$  and  $n$  is large. Then, we use the integral representation of the  $\Gamma$  function, exchange summation and integration, and obtain

$$\Delta E_R(\kappa) = -C \int_0^{\infty} \sum_{\nu=n}^{\infty} \left[ \frac{-\kappa t^{m-1}}{D} \right]^{\nu} t^{K-1/2} e^{-t} dt. \quad (20)$$

The geometric series in this equation can be summed in the Borel sense

$$\Delta E_R(\kappa) = (-1)^{n+1} C (\kappa/D)^n \int_0^{\infty} \frac{t^{(m-1)n + K - 1/2} e^{-t} dt}{1 + \kappa t^{m-1}/D}. \quad (21)$$

To find the large-order formula for  $\Gamma_n$ , we compute the coefficients of the Taylor expansion of  $E_R$  with respect to  $\mu = \tau^2 = 1 - \kappa$ ,

$$\Gamma_n = \frac{1}{n!} \left. \frac{d^n E_R(\mu)}{d\mu^n} \right|_{\mu=0}. \quad (22)$$

If we combine Eqs. (19) and (22), we see that the sum  $\sum_{\nu=0}^{n-1} c_{\nu} (1-\mu)^{\nu}$  does not contribute to  $\Gamma_n$ . Consequently, the large-order formula for  $\Gamma_n$  can be obtained by differentiating the remainder of the sum  $\Delta E_R$  only. Thus, if we replace in Eq. (22)  $E_R$  by  $\Delta E_R$ , and interchange in Eq. (21) integration and differentiation with respect to  $\mu = 1 - \kappa$ , we get

$$\Gamma_n = -\frac{C}{D^n} \int_0^{\infty} \frac{t^{(m-1)n + K - 1/2} e^{-t} dt}{(1 + t^{m-1}/D)^{n+1}}. \quad (23)$$

We derive an analytic large-order formula for  $\Gamma_n$  by constructing an asymptotic approximation according to the Laplace method [13]. We have applied this method for different values of  $m$  and  $K$  and found

$$\Gamma_n = -CD^{(K+1/2)/(m-1)} I_n, \quad (24)$$

where

$$I_n = J_{m,K} \frac{e^{-[m/(m-1)][D(m-1)n]^{1/m}}}{n^{(m-K-1)/m}}, \quad (25)$$

$$J_{2,K} = \frac{e\sqrt{\pi}}{D^{K/2}}, \quad (26)$$

$$J_{m,K} = \frac{\sqrt{2\pi} D^{(m/2 - K - 1)/[m(m-1)]}}{(m-1)^{(m-K-1)/m} m^{1/2}}, \quad m \geq 3. \quad (27)$$

TABLE I. Comparison of the numerical values of the coefficients  $\Gamma_n$  with their values  $\Gamma_n^{\text{LO}}$  given by the large-order formula (28) for the ground state ( $K=0$ ) of the quartic, sextic, octic, and decadic oscillators ( $m=2,3,4,5$ ). For the asymptotic analysis of these data, see Table III. Numbers in square brackets are powers of 10.

$n$	Quartic oscillator		Sextic oscillator		Octic oscillator		Decadic oscillator	
	$\Gamma_n$	$\Gamma_n^{\text{LO}}$	$\Gamma_n$	$\Gamma_n^{\text{LO}}$	$\Gamma_n$	$\Gamma_n^{\text{LO}}$	$\Gamma_n$	$\Gamma_n^{\text{LO}}$
10	-0.88203[-5]	-0.12314[-4]	-0.63983[-4]	-0.18200[-3]	-0.10130[-3]	-0.21369[-3]	-0.10402[-3]	-0.19836[-3]
25	-0.35367[-7]	-0.43060[-7]	-0.28977[-5]	-0.61652[-5]	-0.99835[-5]	-0.16084[-4]	-0.14801[-4]	-0.21856[-4]
50	-0.76355[-10]	-0.86997[-10]	-0.13891[-6]	-0.25088[-6]	-0.11772[-5]	-0.16529[-5]	-0.25970[-5]	-0.33955[-5]
75	-0.71513[-12]	-0.79322[-12]	-0.16773[-7]	-0.28016[-7]	-0.28464[-6]	-0.37581[-6]	-0.84276[-6]	-0.10457[-5]
100	-0.14217[-13]	-0.15531[-13]	-0.31476[-8]	-0.50082[-8]	-0.95738[-7]	-0.12185[-6]	-0.36076[-6]	-0.43422[-6]
125	-0.45589[-15]	-0.49297[-15]	-0.77117[-9]	-0.11855[-8]	-0.39115[-7]	-0.48549[-7]	-0.18138[-6]	-0.21388[-6]

By substituting the expressions for  $C$ ,  $D$ , and  $I_n$  into Eq. (24), we get after some manipulation a remarkably simple large-order formula

$$\Gamma_n = -\frac{2^{K+3/2}a^{K+1/2}b}{\pi K! \sqrt{m}} (bn)^{[(K+1)/m-1]} e^{-\{[m/(m-1)](bn)^{1/m}\}}, \quad (28)$$

$$b = (m-1)D = 4(m-1)B_m a^{1-m}, \quad m \geq 3. \quad (29)$$

For  $m=2$ , the right-hand side of Eq. (28) has to be divided by  $e^2$ , which yields the leading term of Eq. (14).

Analogously to Eq. (14), we assume that the general large-order behavior of  $\Gamma_n$  can be described by the series

$$\Gamma_n = -\frac{2^{K+3/2}a^{K+1/2}b}{\pi K! \sqrt{m}} (bn)^{[(K+1)/m-1]} e^{-\{[m/(m-1)](bn)^{1/m}\}} \times \left( 1 + \sum_{\nu=1}^{\infty} \frac{a_{\nu}^{(K,m)}}{(bn)^{\nu/m}} \right), \quad m \geq 3 \quad (30)$$

where  $a_{\nu}^{(K,m)}$  are expansion coefficients. For  $m=2$ , the right-hand side of Eq. (30) has to be divided by  $e^2$ . Four expansion coefficients  $a_{\nu}^{(K,m)}$  are known for the ground state of the quartic oscillator ( $K=0, m=2, \nu=1, \dots, 4$ ), when they equal the coefficients  $a_{\nu}^{(0)}$  given in [12]. The calculation of the  $a_{\nu}^{(K,m)}$  coefficients for the sextic and higher-order oscillators is more involved and leads to transcendental expressions. We plan to publish results for these oscillators in the future [14]. The series (30) is expected to be only asymptotic.

TABLE II. Comparison of the numerical values of the coefficients  $\Gamma_n$  with their values  $\Gamma_n^{\text{LO}}$  given by the large-order formula (28) for the first excited state ( $K=1$ ) of the quartic, sextic, octic, and decadic oscillators ( $m=2,3,4,5$ ). For the asymptotic analysis of these data, see Table III. Numbers in square brackets are powers of 10.

$n$	Quartic oscillator		Sextic oscillator		Octic oscillator		Decadic oscillator	
	$\Gamma_n$	$\Gamma_n^{\text{LO}}$	$\Gamma_n$	$\Gamma_n^{\text{LO}}$	$\Gamma_n$	$\Gamma_n^{\text{LO}}$	$\Gamma_n$	$\Gamma_n^{\text{LO}}$
10	-0.25353[-3]	-0.66088[-3]	-0.12410[-2]	-0.47993[-2]	-0.16890[-2]	-0.44625[-2]	-0.16329[-2]	-0.37599[-2]
25	-0.19383[-5]	-0.36537[-5]	-0.80553[-4]	-0.22064[-3]	-0.21047[-3]	-0.42235[-3]	-0.27466[-3]	-0.49760[-3]
50	-0.69940[-8]	-0.10439[-7]	-0.52767[-5]	-0.11312[-4]	-0.30768[-4]	-0.51616[-4]	-0.56558[-4]	-0.88795[-4]
75	-0.86211[-10]	-0.11658[-9]	-0.76277[-6]	-0.14460[-5]	-0.84682[-5]	-0.12987[-4]	-0.20266[-4]	-0.29658[-4]
100	-0.20541[-11]	-0.26357[-11]	-0.16180[-6]	-0.28451[-6]	-0.31195[-5]	-0.45252[-5]	-0.93129[-5]	-0.13044[-4]
125	-0.75313[-13]	-0.93534[-13]	-0.43444[-7]	-0.72553[-7]	-0.13661[-5]	-0.19063[-5]	-0.49460[-5]	-0.67184[-5]

We note at the end of our calculations that the integrals in Eqs. (21) and (23) for  $m=2$  can be expressed in terms of the Kummer function  $U(a,b,z)$  [15],

$$\Delta E_R(\kappa) = (-1)^{n+1} C(D/\kappa)^{K+1/2} \Gamma(n+k+1/2) \times U(n+K+1/2, n+K+1/2, D/\kappa), \quad (31)$$

$$\Gamma_n = -CD^{K+1/2} \Gamma(n+K+1/2) U(n+K+1/2, K+1/2, D). \quad (32)$$

### III. NUMERICAL RESULTS

For the quartic oscillator ( $m=2$ ), the large-order formula (28) was tested in [12]. It was shown that this formula gives numerical values of the  $\Gamma_n$  coefficients close to the exact ones starting from  $n$  about 100. Qualitatively, it can be used from  $n$  about 10.

Because of the  $(bn)^{1/m}$  dependence in Eqs. (28) and (30), slower convergence of the large-order formulas to the actual values of  $\Gamma_n$  can be expected with increasing  $m$ . To clarify this question we performed numerical calculation of the  $\Gamma_n$  coefficients for the ground and first excited state ( $K=0,1$ ) of the sextic, octic, and decadic oscillators ( $m=3,4,5$ ). These results are compared with the results for the quartic oscillator in Tables I–III.

The coefficients  $\Gamma_n$  for  $n=0, \dots, 125$  were calculated by the method described in [16]. Numerical values of a few selected  $\Gamma_n$  coefficients are compared with the large-order formula (28) for  $m=2,3,4,5$  and  $K=0,1$  in Tables I and II. The agreement of the numerical values of  $\Gamma_n$  with the large-order formula (28) goes down with increasing  $m$  and  $K$ . We see, however, that even for the first excited state of the dec-

TABLE III. Asymptotic analysis of the ratio  $\Gamma_n/\Gamma_n^{\text{LO}}$  for the ground and first excited state ( $K=0,1$ ) of the quartic, sextic, octic, and decadic oscillators ( $m=2,3,4,5$ ). The leading term  $a_0$  of the asymptotic expansion (33) corresponding to  $n \rightarrow \infty$  was obtained by means of the diagonal Padé approximants based on the values  $\Gamma_n/\Gamma_n^{\text{LO}}$  for  $n=115, \dots, 125$ .

$m$	Ground state	First excited state
	$a_0$	$a_0$
2	1.000037	1.003079
3	1.000747	1.021083
4	1.008496	0.984731
5	0.993720	0.978293

adic oscillator ( $m=5$  and  $K=1$ ) these values agree relatively well. We note that the absolute values of the  $\Gamma_n$  coefficients obtained from the large-order formula (28) are larger than the actual values of  $\Gamma_n$ . We have studied also the ratio appearing in Eq. (30),

$$\Gamma_n/\Gamma_n^{\text{LO}} = a_0 + \frac{a_1^{(K,m)}}{(bn)^{1/m}} + \frac{a_2^{(K,m)}}{(bn)^{2/m}} + \dots, \quad (33)$$

where  $\Gamma_n$  are numerical values of the expansion coefficients and  $\Gamma_n^{\text{LO}}$  are given by the large-order formula (28). For very large  $n$ , this ratio should converge to  $a_0=1$ . Using the Thiele extrapolation built in MAPLE, we extrapolated  $\Gamma_n/\Gamma_n^{\text{LO}}$  to  $n \rightarrow \infty$  and obtained values of  $a_0$  shown in Table III. These values are very close to one and confirm correctness of Eq. (28).

Similarly to [12], we have tested also the validity of the summation rule

$$\Sigma_0 = \sum_{n=0}^{\infty} \Gamma_n = 2K+1. \quad (34)$$

First we calculated the partial sum  $\Sigma_0^N = \sum_{n=0}^N \Gamma_n$  for  $N=125$  from the numerical values of  $\Gamma_n$ . The remaining part of the sum  $\Sigma_0^{\text{LO}} = \sum_{n=N+1}^{\infty} \Gamma_n$  was calculated numerically with the use of Eq. (28). The infinite limit in this sum was replaced by 5000. Results are shown in Table IV. Since we use only the leading term of Eq. (30) here, the summation rules are obeyed with lower accuracy than in the case of the ground state of the quartic oscillator for which four analytic

TABLE IV. Summation rules  $\Sigma_0^N$  and  $\Sigma_0^N + \Sigma_0^{\text{LO}}$  in comparison with the exact value of the summation rule  $\Sigma_0 = 2K+1$  for the coefficients  $\Gamma_n$  of the strong-coupling expansion for the ground and first excited state ( $K=0,1$ ) of the quartic, sextic, octic, and decadic oscillators ( $m=2,3,4,5$ ).  $N=125$ . Numbers in square brackets are powers of 10.

$m$	Ground state		First excited state	
	$\Sigma_0^N - \Sigma_0$	$\Sigma_0^N + \Sigma_0^{\text{LO}} - \Sigma_0$	$\Sigma_0^N - \Sigma_0$	$\Sigma_0^N + \Sigma_0^{\text{LO}} - \Sigma_0$
2	0.339[-14]	-0.265[-15]	0.582[-12]	-0.134[-12]
3	0.160[-7]	-0.805[-8]	0.965[-6]	-0.592[-6]
4	0.142[-5]	-0.298[-6]	0.543[-4]	-0.183[-4]
5	0.951[-5]	-0.139[-5]	0.286[-3]	-0.849[-4]

coefficients  $a_\nu^{0,2}, \nu=1, \dots, 4$  were used [12]. It is seen that the summation rule is obeyed with very good accuracy for  $m=2$  and  $K=0$ . With increasing  $m$  and  $K$ , the accuracy goes down. We see, however, that the summation rule is obeyed with reasonable accuracy even for the first excited state of the decadic oscillator ( $m=5$  and  $K=1$ ). In any case, the inclusion of the large-order formula (28) improves the accuracy of the rule. We note that  $\Sigma_0^N + \Sigma_0^{\text{LO}} - \Sigma_0$  is always negative. It confirms again that the absolute value of  $\Gamma_n$  given by the large-order formula (28) is larger than the absolute value of the actual coefficients  $\Gamma_n$ .

#### IV. CONCLUSIONS

The results of this paper may be summarized as follows: Starting from known large-order behavior of the divergent weak-coupling expansion coefficients  $b_n$ , we derived the general large-order formula (28) for the strong-coupling expansion coefficients  $\Gamma_n$ . In contrast to [12], where the form of this formula was for  $m=2$  conjectured from numerical analysis, Eq. (28) was derived analytically. The large-order formula (28) is very simple and holds for all anharmonicities  $x^{2m}$  with  $m \geq 2$ , and for all states  $K \geq 0$ . We suggested also the more general series (30) which is expected to be only asymptotic. The numerical tests of Eq. (28) were performed for the quartic, sextic, octic, and decadic oscillator. Our results show that the absolute values of the  $\Gamma_n$  coefficients obtained from Eq. (28) are upper bounds to the absolute values of actual  $\Gamma_n$  (see also [12]). This result and the large-order formula (28) show that the renormalized strong-coupling expansion (13) converges for all  $\tau \in [0,1)$ , for all anharmonicities  $x^{2m}$  with  $m \geq 2$ , and for all states  $K \geq 0$ . Therefore the energy  $E(\beta)$  can for all physically relevant  $\beta \in [0, \infty)$  be computed via the *convergent* renormalized strong-coupling expansion (13).

It is remarkable that starting from the large-order behavior (16) of the divergent series (12) it is possible to derive the large-order behavior (28) of the convergent series (13). This can be understood as follows. The weak-coupling expansion  $E_R(\kappa) = \sum_n c_n \kappa^n$  is expanded at the singular point  $\kappa=0$  and, therefore, diverges for any  $\kappa \in (0,1)$ . From the physical point of view, this point is singular since the Hamiltonian  $H_R = p^2 + x^2 + \kappa(x^4/3 - x^2)$  does not have bound states for any  $\kappa < 0$  and the energy  $E_R(\kappa)$  is not analytic at the point  $\kappa=0$ . In the strong-coupling case when  $E_R(\kappa) = \sum_n \Gamma_n (1-\kappa)^n$ , the Hamiltonian  $H_R = p^2 + x^4/3 + (1-\kappa)(x^2 - x^4/3)$  becomes for  $1-\kappa < 0$  the Hamiltonian of the double-well problem which has bound states and the energy  $E_R(\kappa)$  can be, at the point  $\kappa=1$ , analytic. From this point of view, our derivation of Eq. (28) from Eq. (16) is nothing but transformation from the singular point  $\kappa=0$  to the physically more reasonable point  $\kappa=1$ . We plan to publish a more detailed discussion in the future.

The large-order formula (28) has been derived from the formula for the coefficients  $b_n$  which is of semiclassical (JWKB) character. Therefore our formula (28) has the same semiclassical character. In addition, in Eq. (30) we consider the higher-order corrections [17].

The results of this paper can be applied also to more general oscillators such as, for example, to one-dimensional oscillators which have in addition to the  $x^2$  and  $x^{2m}$  terms other

even powers in the potential [7] as well as for isotropic multidimensional problems considered in [18].

Our results show that the renormalized strong-coupling expansion (13) is both from the physical and mathematical point of view the most advantageous perturbative approach to the anharmonic oscillators. Since the perturbation theory of anharmonic oscillators is important not only as model systems in quantum mechanics and quantum field theory but

also in many applications [19], we believe that our results are of some interest.

#### ACKNOWLEDGMENTS

This work was supported in part by the GACR, the GAUK (L.S. and J.Z.), the NSERC (J.C.), the FRVS, and the FCI of Germany (E.J.W.).

- 
- [1] C.M. Bender and T.T. Wu, *Phys. Rev.* **184**, 1231 (1969).
- [2] B. Simon, *Ann. Phys. (N.Y.)* **58**, 76 (1970); S. Graffi, V. Grecchi, and B. Simon, *Phys. Lett.* **32B**, 631 (1970). See also A.V. Turbiner and A.G. Ushveridze, *J. Math. Phys.* **29**, 2053 (1988); A.G. Ushveridze, *Bulg. J. Phys.* **16**, 137 (1989); R. Guardiola, M.A. Solís, and J. Ros, *Nuovo Cimento B* **107**, 713 (1992); F.M. Fernández, *Phys. Lett. A* **166**, 173 (1992); F.M. Fernández and R. Guardiola, *J. Phys. A* **26**, 7169 (1993); W. Janke and H. Kleinert, *Phys. Rev. Lett.* **75**, 2787 (1995).
- [3] C.M. Bender and T.T. Wu, *Phys. Rev. Lett.* **27**, 461 (1971); *Phys. Rev. D* **7**, 1620 (1973).
- [4] Proceedings of the Sanibel Workshop on Perturbation Theory at Large Order, Sanibel Conference, Florida, 1981, edited by P.O. Löwdin and Y. Öhrn [*Int. J. Quantum Chem.* **21**, 1 (1982)] (namely, articles by B. Simon and C.M. Bender).
- [5] *Large-Order Behaviour of Perturbation Theory*, edited by J.C. Le Guillou and J. Zinn-Justin (North-Holland, Amsterdam, 1990).
- [6] M. Reed and B. Simon, *Methods of Modern Mathematical Physics IV: Analysis of Operators* (Academic Press, New York, 1978).
- [7] T.I. Banks and C.M. Bender, *J. Math. Phys.* **13**, 1320 (1972); C.M. Bender and T.T. Wu, *Phys. Rev. D* **7**, 1620 (1973).
- [8] E.J. Weniger, *Phys. Rev. Lett.* **77**, 2859 (1996).
- [9] J. Čížek and E. R. Vrscaj, *Int. J. Quantum Chem., Quantum Chem. Symp.* **20**, 65 (1986); F. Vinette and J. Čížek, *J. Math. Phys.* **32**, 3392 (1991).
- [10] E.J. Weniger, J. Čížek, and F. Vinette, *J. Math. Phys.* **34**, 571 (1993).
- [11] E.J. Weniger, *Ann. Phys. (N.Y.)* **246**, 133 (1996).
- [12] L. Skála, J. Čížek, V. Kapsa, and E.J. Weniger, *Phys. Rev. A* **56**, 4471 (1997).
- [13] C.M. Bender and S.A. Orszag, *Advanced Mathematical Methods for Scientists and Engineers* (McGraw-Hill, New York, 1978).
- [14] J. Zamastil, J. Čížek, and L. Skála (unpublished).
- [15] *Handbook of Mathematical Functions*, edited by M. Abramowitz and I.E. Stegun (Dover Publications, New York, 1965).
- [16] L. Skála and J. Čížek, *J. Phys. A* **29**, L129 (1996); **29**, 6467 (1996).
- [17] H.J. Silverstone, J.G. Harris, J. Čížek, and J. Paldus, *Phys. Rev. A* **32**, 1965 (1985).
- [18] For example, H. Silverstone, E. Harrell, and C. Grot, *Phys. Rev. A* **24**, 1925 (1981); E. Brézin, J.C. Le Guillou, and J. Zinn-Justin, *Phys. Rev. D* **15**, 1544 (1977); S. Hikami and E. Brézin, *J. Phys. A* **12**, 759 (1979).
- [19] J. Čížek, V. Špirko, and O. Bludsky, *J. Chem. Phys.* **99**, 7331 (1993); V. Špirko and J. Čížek, *ibid.* **102**, 8906 (1995); V. Špirko, J. Čížek, and L. Skála, *ibid.* **102**, 8916 (1995); J. Čížek and V. Špirko, *ibid.* **106**, 10 248 (1997).